

PSTricks

pst-optic

Lenses and Mirrors – examples; v.1.00

February 11, 2009

Documentation by
Herbert Voß

Package author(s):
Manuel Luque
Herbert Voß

Contents

I. Lenses	4
1. A simple colored System	4
2. A Magnifier	5
3. Two Lenses	6
4. Real Image	7
5. Virtual Image	8
6. A Microscope	9
7. Telescope	10
8. Lightspeed measured by Foucault	11
 II. Mirrors	 12
9. High Beam Light	12
10 Low Beam Light	13
 III. Refraction	 14
11 Vertical Medium	14
12 Horizontal Medium	15
13 Parallel Rays and a sloping medium	16
14 A Prisma	17
15 A Prisma for Dispersion	17
16 Refraction with different Angles	19
17 Great difference in the Refractionsnumbers	20
18 Total Reflection	21
19 Total Reflection with a sloping medium	22
 IV. Spherical Optic	 23
20 Refraction at a Spherical Surface	23
20.1 Simple Example	23


20.2 Height of an Image	24
21.Thin Convergent Lenses	25
22.Thick Convergent Lenses	26
23.Thin Divergent Lenses	27
24.Thick Divergent Lenses	28
25.\mirrorCVG	29
26.\mirrorDVG	30

Part I.

Lenses

1. A simple colored System

- $\overline{AB} = 2 \text{ cm}$
- $\overline{OA} = -10 \text{ cm}$
- $\overline{OF'} = 3,333 \text{ cm}$
- $\overline{XO} = 2 \text{ cm}$




```

1\begin{pspicture}(-8.5,-3)(8.5,3)
2\rput(0,0){\lens[focus=3.333,0A=-10,AB=2,X0=2,xLeft=-8.5,xRight=8.5,rayColor=red]}
3\pnode(!X0 2.5){L1} \pnode(!X0 -2.5){L2}
4\psOutLine[length=2](L1)(B')\psBeforeLine[length=2](B')(L2){START}
5\pspolygon[style=rayuresJaunes ,linestyle=none](B)(L1)(END)(START)(L2)
6\rput(0,0){\lens[focus=3.333,0A=-10,AB=2,X0=2,xLeft=-8.5,xRight=8.5,rayColor=red,arrowsize=0.2]}
7\end{pspicture}

```

2. A Magnifier




```

1 \begin{pspicture}(-8,-5)(8,3)
2 \rput(0,0){\lens[lensGlass=true,lensWidth=0.4, focus=4,AB=0.5,OA=-2.8,X0=2,drawing=false]
3   \psline[linewidth=0.5pt](xLeft)(xRight)}
4 \pnode(!X0 2.5){L1} \pnode(!X0 -2.5){L2}
5 \psOutLine[length=5.5,linestyle=none](B')(L1){END1}
6 \psBeforeLine[length=6,linestyle=none](L2)(B') {START}
7 \pspolygon[style=rayuresJaunes,linestyle=none](B)(L1)(END1)(START)(L2)
8 \psline[linewidth=1.5\pslinewidth,arrowinset=0]{->}(A)(B)
9 \uput[270](A){A} \uput[90](B){B}
10 \psline[linewidth=1.5\pslinewidth,arrowinset=0,linestyle=dashed]{->}(A')(B')
11 \uput[270](A'){$\mathbf{A'}$} \uput[90](B'){$\mathbf{B'}$}
12 \psset{linecolor=red,arrowsize=0.2}
13 \pcline[nodesepB=-4](B)(0)% Mittelpunktstrahl
14 \psline[linecolor=red,linestyle=dashed](B)(B')% rückwärtige Verlängerung
15 \Arrows(B)(0)% Mittelpunktstrahl
16 \psOutLine[length=2,arrows=->](B)(0){END6}% Mittelpunktstrahl
17 \psline(B)(I)(F')\psOutLine(I)(F'){END2}\Arrows(I)(F')\Arrows(B)(I)
18 \psOutLine[length=1,linestyle=dashed](I')(B') {END3}
19 \psline[linestyle=dashed](B)(F)\psline(B)(I')\Arrows[arrows=->](B)(I')
20 \psline[linestyle=dashed](B')(I')\psline[linestyle=dashed](B')(I)
21 \psOutLine[length=2,arrows=->>](B')(I') {END4}\psOutLine[length=4](B')(I') {END5}
22 \rput(8,0){\psset{linecolor=black}\eye}
23 \end{pspicture}

```

3. Two Lenses

This is a simple system with two lenses, where the `\lens` macro is used only once. The second lens (the left one) is drawn by the `\psline` macro.




```

1 \begin{pspicture}(-8,-5)(8,3)
2 \rput(0,0){\lens[lensScale=0.6,drawing=false,focus=1.5,OA=-1,X0=5,nameF={},nameFi={},AB=-1]
3   \psline[linewidth=1pt](xLeft)(xRight)} %image intermediaire A1B1 au foyer F'1
4 \psline{->}(4,0)(4,-1) %lentille 2
5 % \psline[linewidth=2\pslinewidth, linecolor=blue]{<->}(5,1.5)(5,-1.5)
6 %On place les points essentiels
7 \pnode(-6,0){O1} \pnode(-6,2.5){E1L1} \pnode(-6,-2.5){E2L1}
8 \pnode(4,0){A1} \pnode(4,-1){B1}
9 \rayInterLens(O1)(B1){5}{Inter1L2} %intersection de O1 avec la lentille L2
10 \pcline[nodesepB=-2](Inter1L2)(O1)%rayon venant de l'infini jusqu'e la lentille L2
11 \Parallel(B1)(O1){B1infy}%rayon parallele au precedent et passant par E1L1
12 \Parallel(B1)(O1){B2infy}%rayon passant par E2L2
13 %intersection de la droite passant par E1L1 et B1 avec la lentille L2
14 \rayInterLens(E1L1)(B1){5}{InterE1B1L2}\psline(E1L1)(InterE1B1L2)
15 %intersection de la droite passant par E2L2 et B1 avec la lentille L2
16 \rayInterLens(E2L1)(B1){5}{InterE2B1L2}
17 \psline(E2L1)(InterE2B1L2)
18 \psline[linestyle=dashed]{>}(A')(B')\psline[linestyle=dashed](InterE1B1L2)(B')
19 \psline[linestyle=dashed](InterE2B1L2)(B')\psline[linestyle=dotted](B')(0)
20 \psOutLine[length=3](B')(InterE1B1L2){END}\psBeforeLine[length=3](InterE2B1L2)(B'){START}
21 \pspolygon[style=rayuresJaunes ,linestyle=none](B1infy)(E1L1)(InterE1B1L2)%
22 (END)(START)(InterE2B1L2)(E2L1)(B2infy)
23 \uput[90](A'){$\mathbf{A'}$}\uput[270](B'){$\mathbf{B'}$}
24 \uput[90](A1){$\mathbf{A_1}$}\uput[270](B1){$\mathbf{B_1}$}
25 \uput[225](O1){$\mathbf{O1}$}\uput[45](0){$\mathbf{O2}$}\uput[90](F){$\mathbf{F_2}$}
26 \uput{0.4}[150](F'){$\mathbf{F'_2}$}\uput{0.6}[90](A1){$\mathbf{F'_1}$}
27 \psline[linecolor=red](B1infy)(E1L1)(InterE1B1L2)(END)
28 \psline[linecolor=red](B2infy)(E2L1)(InterE2B1L2)(START)
29 \rput(8,0){\eye}
30 \psline[linewidth=2\pslinewidth, linecolor=blue,arrowsize=0.2,arrowinset=0.5]{<->}(-6,-2.5)(-6,2.5)
31 \end{pspicture}


```

4. Real Image



```
1 \begin{pspicture*}(-7.5,-3)(7.5,3)
2 \rput(0,0){\lens[lensGlass=true,lensWidth=0.5,lensType=DVG,X0=0,AB=2,0A=-4,focus=-6,spotAi=270,
   spotBi=90]%
3 \psline[linewidth=1pt](xLeft)(xRight)
4 \psline[linecolor=red,linestyle=dashed](I')(F)% Verlaengerung des Brennstrahls
5 \psOutLine[length=7](B')(I){END}\psBeforeLine[length=7](I')(B'){START}% permet de definir START
6 \pspolygon[style=rayuresJaunes,linestyle=none](B)(I)(END)(START)(I')
7 \psline(B)(I)(END) \psline(B)(I')(START)
8 \end{pspicture*}
```

5. Virtual Image




```

1 \begin{pspicture*}(-7.5,-6.5)(7.5,7.5)
2 \rput(0,0){\lens[lensType=DVG,lensWidth=0.75,lensHeight=7,focus=-2,0A=-6,AB=4,X0=-1,lensGlass=true
3   ,
4   rayColor=red,yBottom=-5,yTop=5,drawing=false]
5   \psline[linewidth=lpt](xLeft)(xRight)}
6 \pnode(!X0 2.9){L1} \pnode(!X0 -2.5){L2}
7 { \psset{length=4,linestyle=none}
8   \psOutLine(B')(L1){A1} \psOutLine(B')(L2){A2}
9   \pspolygon[style=rayuresJaunes,linestyle=none](B)(L1)(A1)(A2)(L2)
10 \psset{linecolor=red,linestyle=solid}
11 \psline(B)(L1)(A1) \psline(B)(L2)(A2) \psline[linestyle=dashed](B')(L1)
12 \psline[linestyle=dashed](B')(L2) }
13 \psline[linestyle=dashed]{->}(A)(B) \psline{->}(A')(B')
14 \uput[90](B){B}\uput[90](B'){$\mathbf{\mathit{B'}}$}\uput[270](A){A}\uput[270](A'){$\mathbf{\mathit{A'}}$}
15 \end{pspicture*}

```

6. A Microscope




```


1 \begin{pspicture}(-7.5,-5.5)(7.5,3)
2 \rput(0,0){\lens[focus=1.5,0A=-2,AB=0.5,X0=-5,lensGlass=true,lensWidth=0.4,
3   yBottom=-4,yTop=4,drawing=false,lensScale=0.4,nameF=F_1,nameFi=F'_1]
4   \psline[linewidth=1pt](xLeft)(xRight)}
5 \pnode(! X0 1){UPlens1} \pnode(! X0 -1){DOWNlens1}
6 \Transform
7 \rput(0,0){\lens[focus=2,X0=3,lensGlass=true,lensWidth=0.4,yBottom=-4,yTop=4,drawing=false,
8   nameF=F_2,nameFi=F'_2,spotF=90,spotFi=90]}
9 \psline{->}(A1)(B1)\psline{->}(A'1)(B'1)\uput[270](A1){A}\uput[90](B1){B}
10 \uput[270](B'1){$\mathbf{B\_1}$}\uput{0.7}[90](A'1){$\mathbf{A\_1}$}
11 {\psset{linecolor=red}
12 \rayInterLens(I11)(B'1){3}{Inter1L2}\rayInterLens(B1)(O1){3}{Inter2L2}
13 \rayInterLens(UPlens1)(B'1){3}{Inter3L2}\rayInterLens(DOWNlens1)(B'1){3}{Inter4L2}
14 \psline(B1)(I11)(B'1)(Inter1L2)\psline(B1)(Inter2L2)\psline(B1)(UPlens1)(Inter3L2)
15 \psline(B1)(DOWNlens1)(Inter4L2)
16 \psset{length=5}
17 \Parallel(B'1)(0)(Inter3L2){B1inftyRigth}\Parallel(B'1)(0)(Inter4L2){B2inftyRigth}
18 \Parallel(B'1)(0)(Inter2L2){B3inftyRigth}\Parallel(B'1)(0)(Inter1L2){B3inftyRigth}
19 {\psset{length=-5,linestyle=dashed}
20 \Parallel(B'1)(0)(Inter3L2){B1inftyLeft}\Parallel(B'1)(0)(Inter4L2){B2inftyLeft}
21 \Parallel(B'1)(0)(Inter2L2){B3inftyLeft}\Parallel(B'1)(0)(Inter1L2){B3inftyLeft}
22 \pcpline[nodesep=6](B'1)(0)}
23 \pspolygon[style=rayuresJaunes,linestyle=none](B1)(UPlens1)(Inter3L2)%
24   (B1inftyRigth)(B2inftyRigth)(Inter4L2)(DOWNlens1)
25 \psline(B1)(UPlens1)(Inter3L2)(B1inftyRigth)\psline(B2inftyRigth)(Inter4L2)(DOWNlens1)(B1)}
26 \rput(7,0){\eye}
27 \end{pspicture}%

```

7. Telescope




```
\telescope[mirrorFocus=10,posMirrorTwo=8,yBottom=-8]
```



```
\telescope[mirrorFocus=6,posMirrorTwo=5,yBottom=-5]
```

8. Lightspeed measured by Foucault

1849 Foucault (1819-1868) determines with the following configuration the speed of the light.



```

1 \begin{pspicture}(-8,-3.2)(7,4.5)
2 \rput(0,0){\lens[lensWidth=1,lensGlass=true,lensHeight=6,focus=4,drawing=false,AB=2.5]}
3 {\psset{linewidth=0.5pt,linestyle=dashed,arrowsize=5pt,arrows=|<-|}}
4 \psline(-8,0)(4,0)\pcline(-7.75,-3)(0,-3)\lput*{:U}{2f}
5 \pcline(0,-3)(4,-3)\lput*{:U}{f}\pcline(7,0)(7,4)\lput*{:U}{f}
6 \pcline(4,5)(5,5)\lput*{:U}{s}\pcline(5.25,2.3)(5.25,2.8)\lput*{:U}{s} }
7 \uput[90](0,3){\Large L}\uput[45](-7.7,3){\Large B}\uput[45](-7.7,-2){\Large E}
8 \uput[270](3,-0.5){\Large D}\uput[-45](4,0){\Large A=F}\uput[270](3,2){\Large S}
9 \uput[90](4,4){\Large Sp}\uput[90](3.5,3.25){\Large B'}\uput[0](6.3,2.25){\Large B''}
10 \uput[-90](6,1.1){\Large M}\psarc[linewidth=0.5pt]{-7.75,2.5}{0.5}{-90}{0}\qdisk(-7.55,2.3){1pt}
11 \rput{210}(F'){\mirrorTwo}
12 {\psset{fillstyle=solid,fillcolor=lightgray}
13 \rput{210}(4,2.5){\psframe(-1,0)(1,0.2)}\psframe(-8,-3)(-7.75,3)
14 \psframe(3,4)(3.8,4.2)\psframe(4.2,4)(5,4.2) }
15 {\psset{linewidth=1pt, linecolor=red,arrows=->,arrowsize=5pt}
16 \arrowLine[linecolor=blue,arrowOffset=-0.2](F')(4,2.5){2}
17 \arrowLine[linecolor=blue,arrowOffset=-0.2](4,2.3)(6,2.3){1}
18 \qdisk(6,2.3){2pt}
19 \psline[linestyle=dashed,arrows=-](F')(5.1,4)\psline[linestyle=dashed,arrows=-](5,2.8)(6,2.8)
20 \arrowLine(4,4)(F'){3}\arrowLine[linecolor=blue,arrowOffset=-0.2](I)(F'){2}
21 \arrowLine(F')(I){2}\arrowLine[linecolor=blue,arrowOffset=-0.3](-7.75,2.5)(I){3}
22 \arrowLine(I)(-7.75,2.5){3}
23 \psframe(5.5,1.1)(6.5,3.5)
24 \multido{\r=1.3+0.2}{12}{\psline(6.1,\r)(6.5,\r)}
25 \end{pspicture}


```

Sp chink;
 D rotating mirror;
 L collecting lens;
 E end mirror;
 S half diaphanous mirror;
 M scale

Part II.

Mirrors

9. High Beam Light




```

1 \begin{pspicture}(-1.5,-5.5)(10,5.5)
2 \rput(0,0){\beamLight[drawing=false,mirrorDepth=4.75,mirrorWidth=0.1,mirrorHeight=10,linecolor=lightgray]}
3 \makeatletter
4 \pst@getcoor{Focus}\pst@tempf
5 \psset{linewidth=1pt,linecolor=red}
6 \multido{\n=60+5}{18}{%
7   \mirrorCVGRay[linecolor=red,mirrorDepth=4.75,mirrorHeight=10,linewidth=1pt](Focus)(!%
8     /XF \pst@tempf pop \pst@number\psxunit div def \n\space cos XF add \n\space sin neg){Endd1}
9   \psOutLine[arrows=->,length=.25](Endd1)(Endd1''){Endd2}%
10  \mirrorCVGRay[linecolor=red,mirrorDepth=4.75,mirrorHeight=10,linewidth=1pt](Focus)(!%
11    /XF \pst@tempf pop \pst@number\psxunit div def \n\space cos XF add \n\space sin ){End1}
12  \psOutLine[arrows=->,length=.25](End1)(End1''){End2}}
13 \makeatletter
14 \end{pspicture}

```

10. Low Beam Light




```
1 \begin{pspicture}(-1.5,-5)(10,5)
2 \rput(0,0){\beamLight[drawing=false,mirrorDepth=4.75,mirrorWidth=0.1,mirrorHeight=10,linecolor=
    lightgray]}
3 \psset{linewidth=1pt,linecolor=red}
4 \multido{\n=70+5}{20}{%
5   \psline(2.75,-0.2)(! \n\space cos 2.75 add \n\space sin )
6   \mirrorCVGRay[linecolor=red,mirrorDepth=4.75,mirrorHeight=10,linewidth=1pt](2.75,-0.2)%
7     (! \n\space cos 2.75 add \n\space sin ){End1}
8   \psOutLine[arrows=->,length=.25](End1)(End1''){End2}}
9 \end{pspicture}
```

Part III.

Refraction

11. Vertical Medium

Refraction numbers are $n_1 = 1$ and $n_2 = 1.5$:




```

1 \begin{pspicture}[showgrid=true](-5,-3)(5,3)
2 \pnode(-1,-2.5){A}\pnode(1,-2.5){B}\pnode(1,2.5){C}\pnode(-1,2.5){D}
3% \rotateFrame(A)(B)(C)(D){10}
4 \uput[-135](A){A}\uput[-45](B){B}\uput[45](C){C}\uput[135](D){D}
5 \pspolygon[fillcolor=lightgray, fillstyle=solid, linecolor=blue](A)(B)(C)(D)
6% 1.
7 \refractionRay(-3,-3)(-2,-2)(D)(A){1}{1.5}{END}
8 \psset{linecolor=red, linewidth=2pt, arrowsize=5pt, arrows=->}
9 \arrowLine(-3,-3)(END){2}\ABinterCD(END')(C)(B){Out}
10 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(C)(B){1.5}{1}{Q}
11 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
12% 2.
13 \refractionRay(-3,0)(-2,0)(A)(D){1}{1.5}{END}
14 \psset{linecolor=green, linewidth=2pt, arrowsize=5pt, arrows=->}
15 \arrowLine(-3,0)(END){2}\ABinterCD(END')(C)(B){Out}
16 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(C)(B){1.5}{1}{Q}
17 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
18% 3.
19 \refractionRay(-3,3)(-2,2)(D)(A){1}{1.5}{END}
20 \psset{linecolor=blue, linewidth=2pt, arrowsize=5pt, arrows=->}
21 \arrowLine(-3,3)(END){2}\ABinterCD(END')(C)(B){Out}
22 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(C)(B){1.5}{1}{Q}
23 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
24 \end{pspicture}

```

12. Horizontal Medium

Refraction numbers are $n_1 = 1$ and $n_2 = 1.5$:




```

1 \begin{pspicture}[showgrid=true](-5,-4)(5,3)
2 \pnode{-2.5,-1}{A}\pnode{2.5,-1}{B}\pnode{2.5,1}{C}\pnode{-2.5,1}{D}
3 \%rotateFrame(A)(B)(C)(D){10}
4 \uput[-135](A){A}\uput[-45](B){B}\uput[45](C){C}\uput[135](D){D}
5 \pspolygon[fillcolor=lightgray,fillstyle=solid,linecolor=blue](A)(B)(C)(D)
6 \% 1.
7 \refractionRay(-3,3)(-2,2)(C)(D){1}{1.5}{END}
8 \psset{linecolor=red,linewidth=2pt,arrowsize=5pt,arrows=->}
9 \arrowLine(-3,3)(END){2}\ABinterCD(END')(B)(A){Out}
10 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(B)(A){1.5}{1}{Q}
11 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
12 \% 2.
13 \refractionRay(0,3)(0,1)(C)(D){1}{1.5}{END}
14 \psset{linecolor=green,linewidth=2pt,arrowsize=5pt,arrows=->}
15 \arrowLine(0,3)(END){2}\ABinterCD(END')(A)(B){Out}
16 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(B)(A){1.5}{1}{Q}
17 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
18 \% 3.
19 \refractionRay(3,3)(2,2)(C)(D){1}{1.5}{END}
20 \psset{linecolor=blue,linewidth=2pt,arrowsize=5pt,arrows=->}
21 \arrowLine(3,3)(END){2}\ABinterCD(END')(B)(A){Out}
22 \arrowLine(END)(Out){1}\refractionRay(END)(Out)(B)(A){1.5}{1}{Q}
23 \arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
24 \end{pspicture}

```

13. Parallel Rays and a sloping medium

Refraction numbers are $n_1 = 1$ and $n_2 = 1.5$:




```

1 \begin{pspicture}[showgrid=true](-7,-1)(5,6)
2 \pnode(0,0){A}\pnode(2,0){B}\pnode(2,5){C}\pnode(0,5){D}
3 \rotateFrame(A)(B)(C)(D){40}
4 \uput[-135](A){A}\uput[-45](B){B}\uput[30](C){C}\uput[135](D){D}
5 \pspolygon[fillcolor=lightgray,fillstyle=solid,linecolor=blue](A)(B)(C)(D)
6 \psset{linecolor=red,linewidth=2pt,arrowsize=5pt,arrows=->}
7 \multido{\r=1.10+0.28}{8}{%
8   \refractionRay(-6.00,\r)(-3.00,\r)(A)(D){1}{1.5}{End}
9   \arrowLine(-6.00,\r)(End){2}\ABinterCD(End)(End')(C)(B){Out}
10 \arrowLine(End)(Out){1}\refractionRay(End)(Out)(C)(B){1.5}{1}{Q}
11 \psline(Q)(Q')\psOutLine[length=3](Q)(Q'){End}}
12 \end{pspicture}

```

14. A Prisma

Refraction numbers are $n_1 = 1$ and $n_2 = 1.5$:



```


1 \begin{pspicture}[showgrid=true](-7,0)(5,6)
2 \pnode(-3,1){A}\pnode(1,1){B}\pnode(-1,5){C}\uput[-135](A){A}\uput[-45](B){B}\uput[30](C){C}
3 \pspolygon[fillcolor=lightgray, fillstyle=solid, linecolor=blue](A)(B)(C)
4 \psset{linecolor=red, linewidth=2pt, arrowsize=5pt, arrows=->}
5 \multido{\rA=0.6+0.2, \rB=1.5+0.2}{7}{%
6   \refractionRay(-6,\rA)(-4,\rB)(C)(A){1}{1.5}{END}\arrowLine(-6,\rA)(END){2}
7   \ABinterCD(END')(C)(B){Out}\arrowLine(END)(Out){1}
8   \refractionRay(END)(Out)(C)(B){1.5}{1}{Q}\psline(Q)(Q')\psOutLine[length=3](Q)(Q'){End}}
9 \end{pspicture}

```

15. A Prisma for Dispersion


The following figure shows the light dispersion with realistic values for the refractions numbers of the different light colors.

	darkblue	bluegreen	yellow	red	darkred
n for glass	1.528	1.523	1.517	1.514	1.511



16. Refraction with different Angles

Refraction numbers are $n_1 = 1$ and $n_2 = 1.5$:




```

1 \begin{pspicture}[showgrid=true](-6,-5)(6,5)
2 \pnode(-6,-1){A}\pnode(6,-1){B}\pnode(6,1){C}\pnode(-6,1){D}
3 \uput[-135](A){A}\uput[-45](B){B}\uput[30](C){C}\uput[135](D){D}
4 \pspolygon[fillcolor=lightgray,fillstyle=solid,linecolor=blue](A)(B)(C)(D)
5 \psline[linewidth=0.5pt](0,-5)(0,5)
6 \psset{linecolor=red,linewidth=1.5pt,arrowsize=5pt,arrows=->}
7 \multido{\n=30+5}{25}{%
8   \refractionRay(5;\n)(0,1)(C)(D){1}{1.5}{END}\arrowLine(5;\n)(END){2}
9   \ABinterCD(END)(END')(B)(A){Out}\arrowLine(END)(Out){1}
10 \refractionRay(END)(Out)(B)(A){1.5}{1}{Q}\psline(Q)(Q')\psOutLine[length=3](Q)(Q'){End}}
11 \end{pspicture}

```

17. Great difference in the Refractionsnumbers

Refractionnumbers are $n_1 = 1$ and $n_2 = 4$:




```

1 \begin{pspicture}[showgrid=true](-5,-1)(2,6)
2 \pnode(0,0){A}\pnode(2,0){B}\pnode(2,5){C}\pnode(0,5){D}\rotateFrame(A)(B)(C)(D){45}
3 \uput[-135](A){A}\uput[-40](B){B}\uput[45](C){C}\uput[135](D){D}
4 \pspolygon[fillcolor=lightgray, fillstyle=solid, linecolor=blue](A)(B)(C)(D)
5 \refractionRay(-2.5,-1)(-2,1)(A)(D){1}{4}{END}
6 \psset{linecolor=red, linewidth=2pt, arrowsize=5pt, arrows=->}
7 \arrowLine(-2.5,-1){2}\ABinterCD(END')(END')(C)(B){Out}\arrowLine(END)(Out){1}
8 \refractionRay(END)(Out)(C)(B){4}{1}{Q}\arrowLine(Q)(Q'){1}\psOutLine[length=2](Q)(Q'){End}
9 \end{pspicture}

```


18. Total Reflection

Refraction numbers are $n_1 = 2$ and $n_2 = 1$:



19. Total Reflection with a sloping medium

Refraction numbers are $n_1 = 2$ and $n_2 = 1$:



```

1 \begin{pspicture}[showgrid=true](-8,-6)(8,6)
2 \pnod(-6,1.5){A}\pnod(6,1.5){B}\pnod(6,-1.5){C}\pnod(-6,-1.5){D}
3 \rotateFrame(A)(B)(C)(D){30}\uput[90](A){A}\uput[135](B){B}\pnod(0,0){START}
4 \pspolygon[fillcolor=lightgray,fillstyle=solid,linecolor=blue](A)(B)(C)(D)
5 \psset{linecolor=red,linewidth=1.5pt,arrowsize=5pt,arrows=->}
6 \multido{\n=50+5}{29}{%
7   \refractionRay(START)(1;\n)(A)(B){2}{1}{END}\arrowLine(START)(END){2}
8   \arrowLine(END)(END')[1]\psOutLine[length=3](END)(END')[Q]\arrowLine(END')(Q){3}}
9 \end{pspicture}


```

Part IV.

Spherical Optic

20. Refraction at a Spherical Surface

20.1. Construction for finding the position of the image point P' of a point object P formed by refraction at a spherical surface




```

1 \begin{pspicture*}[showgrid=true](-10,-4)(3,4)
2 \rput(0,0){\lensSPH[lensType=CVG,lensHeight=12,lensWidth=10,yBottom=-6,yTop=6,xLeft=-6,xRight=6,
3   drawing=false]}
4 \pnode(-9,0){P}\psdots(P)\uput[-90](P){P}\psline(P)(xRight)\lensSPHRay(P)(-5,2){1}{9}{Q}
5 \psline(P)(Q)(Q')\psdots(Q)\uput[90](Q){B}\ABinterCD(Q)(Q')(0,0)(5,0){P'}
6 \psdots(Q')\uput[-90](P'){P'}\psline[lineWidth=0.5pt, linecolor=black](Center')(Q)
7 \psline[lineWidth=0.5pt, linecolor=black](Q)(Q|0,0)\psdots(Center')\uput[-90](Center'){C}
8 \end{pspicture*}

```

20.2. Construction for determining the height of an image formed by refraction at a spherical surface




```

1 \begin{pspicture*}[showgrid=true](-13,-3)(3,5)
2 \rput(0,0){\lensSPH[lensType=CVG,lensHeight=12,lensWidth=10,yBottom=-4,yTop=4,xLeft=-5,xRight=5,
3   drawing=false]}
4 \psset{linecolor=red,linewidth=1.5pt,dotstyle=|}
5 \pnode(-12,0){P}\psdots(P)\uput[-90](P){P}\pnode(-12,4){Q}\psdots(Q)\uput[90](Q){Q}
6 \psline[linecolor=blue,linewidth=3pt,arrows=->](P)(Q)\psline(P)(xRight)
7 \lensSPHRay(Q)(Center')\lensSPHRay(Q)(-5,0)\lensSPHRay(Q)(-5,0)\lensSPHRay(Q)(-5,0)%
8 \psline(Q)(S1')\psline(Q)(S2)(S2')\ABinterCD(Q)(S1')(S2)(S2'){Q'}
9 \pnode(Q'|0,0){P'}\psline[linecolor=blue,linewidth=3pt,arrows=->](P')(Q')
10 \uput[90](P'){P'}\uput[-90](Q'){Q'}\psdots(Center')\uput[90](Center'){C}
11 \end{pspicture*}

```

21. Thin Convergent Lenses

If the two spherical surfaces are close enough we can call such a lens a **thin lens**. The following figure shows the behaviour of such a lens with real rays.




```

1 \psset{xunit=0.75cm}
2 \begin{pspicture*}[showgrid=true](-10,-4)(10,4)
3 \rput(0,0){\lensSPH[lensType=CVG,lensHeight=7,lensWidth=1.25,yBottom=-5,yTop=5,xLeft=-12,xRight
=12,%
4 AB=2,0A=-9,refractA=1,refractB=2,drawing=true,rayColor=red]}
5 \end{pspicture*}

```

22. Thick Convergent Lenses


There is no real image possible.



```
\begin{pspicture}(-10,-4)(10,4)
\rput(0,0){\lensSPH[lensType=CVG,lensHeight=7,lensWidth=2,yBottom=-5,yTop=5,xLeft=-12,xRight=12,%
3 AB=2,0A=-9,refractA=1,refractB=2,drawing=true,rayColor=red]}
\end{pspicture}
```

23. Thin Divergent Lenses


If the two spherical surfaces are close enough we can call such a lens a **thin lens**. The following figure shows the behaviour of such a lens with real rays.



```
1 \begin{pspicture*}[showgrid=true](-6,-3)(6,3)
2 \rput(0,0){\lensSPH[lensType=DVG,lensWidth=0.1,lensDepth=0.2,AB=1,OA=-5,drawing=true,rayColor=red
3 ]}
```


24. Thick Divergent Lenses

There is no real image possible.



```
\begin{pspicture*}[showgrid=true](-6,-3)(6,3)
\put(0,0){\lensSPH[lensType=DVG,lensWidth=1,lensDepth=1,AB=1,OA=-5,drawing=true,rayColor=red]}
\end{pspicture*}
```

25. \mirrorCVG




```

1 \begin{pspicture*}[showgrid=true](-1,-5)(8,5)
2 \rput(0,0){\mirrorCVG[mirrorType=SPH,drawing=false,yBottom=-4,yTop=4,mirrorHeight=8,mirrorDepth=3]
3   \qdisk(Center){2pt}\qdisk(Focus){2pt}
4   \uput[-90](Center){Center}\uput[-90](Focus){F}\psline(0)(xRight)}
5 \multido{\rA=-3.50+0.25}{5}{%
6   \mirrorCVGRay[mirrorType=SPH,linecolor=red](6,\rA)(4,\rA){E}
7   \psOutLine[linecolor=red,length=4](E')(E'')\EEEnd}
8 \multido{\rA=-2.25+0.25}{19}{%
9   \ABinterSPHLens(6,\rA)(4,\rA)(Center){Ptemp}
10  \reflectionRay[mirrorType=SPH](5,\rA)(Ptemp){E}
11  \psline[linecolor=red](6,\rA)(Ptemp)(E)\psOutLine[linecolor=red,length=6](Ptemp)(E)\EEEnd}
12 \multido{\rA=2.50+0.25}{5}{%
13   \mirrorCVGRay[mirrorType=SPH,linecolor=red](6,\rA)(4,\rA){E}
14   \psOutLine[linecolor=red,length=4](E')(E'')\EEEnd}
15 \end{pspicture*}

```

26. \mirrorDVG



```

1 \begin{pspicture*}[showgrid=true](-5,-5)(8,5)
2 \rput(0,0){%
3   \mirrorDVG[mirrorType=SPH,drawing=false,yBottom=-4,yTop=4,mirrorHeight=8,mirrorWidth=0.25,
4     mirrorDepth=2.5]
5   \qdisk(Center){2pt}\qdisk(Focus){2pt}\uput[-90](Center){C}\uput[-90](Focus){F}
6   \psline(xLeft)(xRight)}
7   \multido{\rA=-3.00+0.25}{25}{%
8     \ABinterSPHLens(7,\rA)(4,\rA)(Center){Ptemp}\reflectionRay[mirrorType=SPH](5,\rA)(Ptemp){E}
9     \arrowLine[linecolor=red,linewidth=1.5pt](7,\rA)(Ptemp){1}
10    \psline[linecolor=red,arrows=->,linewidth=1.5pt](Ptemp)(E)
11    \psOutLine[linecolor=red,length=6,linewidth=1.5pt,arrows=->](Ptemp)(E){EEEnd}
12    \psOutLine[linecolor=red,length=3,linestyle=dashed,linewidth=0.5pt](E)(Ptemp){EEEnd}}}
13 \end{pspicture*}

```

Index

A

- AB, 25–28
- \ABinterCD, 14–17, 23, 24
- \ABinterSPHLens, 29, 30
- \arrowLine, 14–17
- \Arrows, 5, 6

B

- \beamLight, 12, 13

C

- CVG, 25, 26

D

- DVG, 27, 28

E

- \eye, 5, 6

K

- Keyword
 - AB, 25–28
 - length, 14
 - lensType, 25–28
 - mirrorDepth, 13
 - mirrorHeight, 13
 - mirrorType, 29, 30
 - refractA, 25, 26
 - refractB, 25, 26

L

- length, 14
- \lens, 5–9
- \lensSPH, 23, 24, 27, 28
- \lensSPHRay, 23, 24
- lensType, 25–28

M

- Macro
 - \ABinterCD, 14–17, 23, 24
 - \ABinterSPHLens, 29, 30
 - \arrowLine, 14–17
 - \Arrows, 5, 6
 - \beamLight, 12, 13
 - \eye, 5, 6
 - \lens, 5–9
 - \lensSPH, 23, 24, 27, 28

- \lensSPHRay, 23, 24
- \mirrorCVG, 29
- \mirrorCVGRay, 12, 13
- \mirrorDVG, 30
- \Parallel, 9
- \polygon, 16
- \psBeforeLine, 4–9
- \psline, 6
- \psOutLine, 4–9, 12, 13, 17
- \rayInterLens, 6, 9
- \refractionRay, 14–17, 29, 30
- \telescope, 10
- \mirrorCVG, 29
- \mirrorCVGRay, 12, 13
- mirrorDepth, 13
- \mirrorDVG, 30
- mirrorHeight, 13
- mirrorType, 29, 30

P

- \Parallel, 9
- \polygon, 16
- \psBeforeLine, 4–9
- \psline, 6
- \psOutLine, 4–9, 12, 13, 17

R

- \rayInterLens, 6, 9
- refractA, 25, 26
- refractB, 25, 26
- \refractionRay, 14–17, 29, 30

S

- SPH, 29, 30

T

- \telescope, 10

V

- Value
 - CVG, 25, 26
 - DVG, 27, 28
 - SPH, 29, 30